




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 237/06, 237/20, A61K 31/16		A1	(11) International Publication Number: WO 98/03472
			(43) International Publication Date: 29 January 1998 (29.01.98)
(21) International Application Number: PCT/EP97/03773 (22) International Filing Date: 15 July 1997 (15.07.97) (30) Priority Data: MI96A001544 23 July 1996 (23.07.96) IT (71) Applicant (for all designated States except US): CHIESI FARMACEUTICI S.P.A. [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): CHIESI, Paolo [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). VENTURA, Paolo [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). DELCANALE, Maurizio [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). DE FANTI, Renato [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). ARMANI, Elisabetta [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). VILLETTI, Gino [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). PIETRA, Claudio [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (74) Agent: MINOJA, Fabrizio; Studio Consulenza Brevettuale, Via Rossini, 8, I-20122 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: ALPHA-AMINO ACID AMIDES, PREPARATION THEREOF AND THE THERAPEUTICAL USE THEREOF

(57) Abstract

The present invention relates to serinamide, glycineamide, alanineamide and phenylalanineamide derivatives of formula (I) wherein R, R', R₁ and R₂ are as defined in the disclosure. The compounds (I) are useful for the treatment of neurological diseases.



FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ALFA-AMINO ACID AMIDES, PREPARATION THEREOF AND THE THERAPEUTICAL USE THEREOF

The present invention relates to α -amino acid amides, processes for the preparation thereof and pharmaceutical compositions containing them.

More precisely, the invention relates to serinamide, glycinamide, alaninamide and phenylalaninamide derivatives of general formula (I):



wherein:

R is straight or branched alkyl; cycloalkylalkyl; arylalkyl or phenylalkyl optionally substituted at the ring with alkyl, halogen or haloalkyl; fused or non-fused aryl optionally substituted with alkyl, alkoxy, halogen or haloalkyl;

R' is hydrogen; alkyl; phenyl; phenylalkyl;

R₁ is optionally acylated C₁-C₄ hydroxyalkyl or phenylalkyl;

R₂ is hydrogen; alkyl; phenyl; phenylalkyl;

with the proviso that when R is aryl, R₁ can be hydrogen or alkyl C₁-C₄.

An alkyl group if not otherwise specified is preferably a C₁-C₁₀ alkyl group such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, 2-ethylpentyl, 1-ethylheptyl, 1-methyloctyl, 4-heptyl.

A cycloalkylalkyl group is preferably a group

2

having 1 to 3 carbon atoms in the alkyl moiety and 3 to 7 carbon atoms in the cycloalkyl moiety such as cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, 1-(5-norbornylenyl)ethyl.

5 An optionally substituted arylalkyl or phenylalkyl group is preferably 2-naphthalenylmethyl, benzyl, phenethyl, phenylpropyl, phenylbutyl, 3-(4-methylphenyl)propyl, 3-(4-fluorophenyl)propyl, 3-(4-chlorophenyl)propyl, 3-(4-trifluoromethylphenyl)propyl,
10 3-phenyl-1-methylpropyl, 2-phenyl-1-methylethyl, 3-phenyl-3-methyl-propyl, 1-phenylethyl.

A fused or non-fused aryl group is preferably 1,2,3,4-tetrahydro-2-naphthalenyl, 2-indanyl optionally substituted by one or more alkoxy, halogen or haloalkyl
15 groups.

An acylated C₁-C₄ hydroxyalkyl group is preferably acetoxyalkyl, propanoyloxyalkyl, 2-methylpropanoyloxyalkyl, benzoyloxyalkyl group.

A class of preferred compounds is that wherein:

20 R₁ is CH₂OH; R is C₃-C₁₀ alkyl, C₂-C₄ phenylalkyl, 1,2,3,4-tetrahydro-2-naphthalenyl or 2-indanyl optionally substituted with alkyl, alkoxy, halogen or haloalkyl; R' is hydrogen or methyl and R₂ is hydrogen or methyl.

25 A particularly preferred sub-class is that wherein R₁ is CH₂OH; R is phenyl-(C₂-C₃)-alkyl or 1,2,3,4-tetrahydro-2-naphthalenyl, or 2-indanyl optionally substituted by one or more alkoxy, halogen, haloalkyl groups; R' and R₂ are hydrogen.

30 A second class of preferred compounds is that wherein R₁ is hydrogen or methyl, R is 1,2,3,4-

3

tetrahydro-2-naphthalenyl or 2-indanyl optionally substituted by one or more alkoxy, halogen, haloalkyl groups and R' and R₂ are hydrogen.

5 The compounds of the invention can be in the form of organic or inorganic acid addition salts.

Moreover they can have one or more asymmetric carbon atoms, therefore they can be used both in the form of mixtures containing more diastereoisomers in any ratio, and in the form of racemic mixtures containing
10 couples of enantiomers in equal or different ratios, and in the form of optically pure compounds.

The compounds of the invention can be used in the treatment of chronic neurodegenerative diseases, such as Alzheimer's disease, various forms of dementia,
15 Parkinson's disease, Huntingdon's disease or acute neurodegenerative impairments such as stroke and head injuries; in the treatment of epilepsy and depression.

PRIOR ART

GB patent 2048852 (Continental Pharma S.A.)
20 discloses 2-aminoacetamide (commonly referred to as glycineamide) derivatives which can be used in the treatment of epilepsy, in the treatment of dyskinesias such as Parkinson's disease, in the treatment of memory disorders and possibly in the treatment of depression.

25 Some of the disclosed compounds, orally administered in doses of 10-100 mg/kg, showed anticonvulsive effects against bicuculline-induced tonic convulsions in mouse.

2-n-Pentylaminoacetamide (in the following referred
30 to with the nonproprietary name milacemide) and its hydrochloride were particularly studied.

Milacemide was used as the reference compound to test the pharmacological activity of the compounds of the present invention.

5 EP-B1-0400495 (Farmitalia Carlo Erba) discloses α -aminocarboxamide N-phenylalkyl substituted derivatives.

Examples of particularly preferred compounds are aminopropionamide (particularly alaninamide and serinamide) and aminoacetamide (glycinamide) N-phenylalkyl substituted derivatives. Said compounds are
10 active on Central Nervous System and can be used as antiepileptics, anti-Parkinsons, antineurodegeneratives, antidepressants, hypnotics and antispastics.

The activity of the compounds has been evaluated in the mouse as the anticonvulsive action against
15 bicuculline- or 3-mercaptopropionic acid- induced convulsions.

The compounds described in EP-B1-0400495 are also potent monoamino oxidase (MAO) inhibitors.

20 WO 94/22808 and WO 94/22809 (Pharmacia/Farmitalia Carlo Erba) disclose other aminopropionamide derivatives, acting on Central Nervous System, respectively arylalkoxybenzyl- and arylalkylaminobenzyl-substituted.

25 One of the most representative compounds of the invention in WO 94/22808 is FCE28245, chemically 2-{4-[3-phenylpropyl]oxybenzyl]-amino-3-hydroxypropanamide methanesulfonate claimed to be active in the test of the electro-shock convulsions in the mouse.

30 PCT n° WO 95/18617 (Teva-Technion) and PCT n. WO 96/21640 (Teva-Lemmon) describe 1-aminobenzocycloalkane derivatives, such as 1-aminoindans and 1-aminotetralins,


5

which can be used in the treatment of Parkinson's disease, dementias, epilepsy and in post-traumatic diseases.


However, some of the described compounds, such as
5 racemic N-(2-acetamido)-1-aminoindan and its optically active forms; N-(2-acetamido)-6-fluoro-1-aminoindan; N-(2-acetamido)-1-aminotetralin; N,N-di-(2-acetamido)-1-aminoindan and N-(2-propionamido)-1-aminoindan were not very effective in the anticonvulsant activity test and
10 don't seem endowed with a particularly favourable therapeutic index.

Now it has been found that α -amino acids amides of general formula (I) are characterized by a higher efficacy and/or a better pharmacological profile than
15 the prior art compounds.


The compounds of the invention can be prepared reacting amino acids esters or amides of formula II



wherein R_1 is as defined above and X is an alkoxy group or a NHR_2 group, wherein R_2 is as defined above, with compounds of formula III



wherein Y is an oxygen atom or a NH group, whereas R_3 and R_4 , which are the same or different, are hydrogen or, together with the carbon atom they are linked to,
30 they form one of the groups R or R_1 , as defined above, to give compounds of formula IV



which can then be transformed into compounds of formula I by means of one or more of the following reactions:

- when X is an alkoxy group, reaction with an amine of formula $\text{R}_2\text{-NH}_2$;
- N-alkylation;
- acylation of any hydroxy group present in R_1 ;
- salification and/or optical resolution.
- elimination of any protecting groups.

A first embodiment of the process described above involves the reductive amination of a compound of formula II or a salt thereof (generally the hydrochloride) wherein X is an alkoxy group, for example methoxy, with a compound of general formula III wherein Y is oxygen and the subsequent reaction with an amine of formula $\text{R}_2\text{-NH}_2$.

The reductive amination is carried out according to conventional methods, using stoichiometric amounts or slight excesses of the reagents, at temperatures ranging from 0 to 40°C and in organic solvents such as alcohols or acetonitrile. A hydride such as sodium cyanoborohydride or hydrogen in the presence of a catalyst such as Pd on-carbon can be used as a reducing agent.

The subsequent amidation reaction is carried out using an amine excess in water or in an organic solvent, particularly methanol or ethanol, at room temperature or heating in a chemical reactor.


A second embodiment involves the reductive

7


amination of a compound II wherein X is a R_2 -NH group, according to the procedures already described.

Finally, a third embodiment involves the transimination of a compound II wherein X is R_2 -NH- with an imine compound (generally a phenylimine) III wherein Y is NH. The reaction is carried out in an organic solvent, for example an alcohol, methylene chloride or acetonitrile at a temperature from 0° to 40°C. The subsequent reduction of the resulting compound is carried out in an organic solvent, generally an alcohol such as ethanol or methanol, using a hydride such as sodium borohydride as reducing agent, at a temperature from 0° to 40°C.

Alternatively, the compounds I can also be prepared by condensation of an alfa-halogen ester of formula V



wherein R_1 is as defined above and preferably H, W is a halogen atom (generally chlorine or bromine) and R_5 is an alkyl group, with an amine of formula VI



wherein R_3 and R_4 are as defined above, and subsequent amidation with the amine R_2 -NH₂. The condensation of compound V with compound VI is carried out in an organic solvent, for example acetonitrile, alcohol, dimethylformamide at a temperature from 40° to 140°C in the presence of an acid-binding agent, for example potassium carbonate and preferably in the presence of

catalytic amounts of potassium iodide. The amidation is then carried out as described above.

It is also possible to condense an alfa-halogen amide with the amine VI.

5 For the envisaged therapeutical uses, compounds I will be formulated in suitable pharmaceutical compositions which are a further object of the invention.

10 Said compositions will typically contain 1 to 1000 mg of active ingredient, particularly 10 to 100 mg, and will be administered one or more times a day depending on the disease, the pharmacokinetics of the selected active ingredient and the conditions (weight, sex, age) of the patient.

15 The compositions will be prepared using conventional techniques and excipients as described for example in Remington's Pharmaceutical Sciences Handbook, Mack. Pub., N.Y., U.S.A., and will be administered by the oral, parenteral or rectal route. Examples of
20 formulations comprise tablets, capsules, syrups, granulates, sterile injectable solutions or suspensions, suppositories and the like.

The following examples further illustrate the invention.

25 **EXAMPLES**

Example 1

a) Preparation of N-(3-phenylpropyl)-L-serine methyl ester hydrochloride

30 L-Serine methyl ester hydrochloride (0.9 moles, 14 g), triethylamine (0.9 moles, 9.1 g) and 3-phenylpropionaldehyde (0.9 moles, 12.1 g) are dissolved

in dry methanol (370 ml) in a Parr bottle and hydrogenated under 45 psi in the presence of 10% Pd/C, until the hydrogen absorption ceases.

5 The catalyst is filtered off and the filtrate is evaporated to dryness under vacuum. The resulting oil is taken up with methylene chloride (500 ml), the organic solution is washed with water and evaporated to dryness under vacuum to obtain a pale yellow oil.

10 The product is recovered as the hydrochloride by dissolution in ethyl ether (800 ml) and acidification with methanol hydrochloric acid. The precipitate is filtered and dried under vacuum at 45°C.

Yield: 17.5 g (71%) - m.p. = 126-129°C

15 b) Preparation of (-)-(S)-3-hydroxy-2-(3-phenylpropyl-amino)propanamide hydrochloride (CHF 2803.01)

The product obtained in a)! (0.06 moles, 17 g) is dissolved in water (500 ml) and alkalized with 10% aqueous potassium carbonate to pH = 8. The free base is extracted with methylene chloride and evaporated to
20 dryness under vacuum. The resulting pale yellow oil (16.3 g) is dissolved in methanol (150 ml). Ammonia is bubbled through the solution, cooled at -5°C, to a ~15M concentration. The hermetically sealed system is reacted for 5 days at room temperature (r.t.), then evaporated
25 to dryness under vacuum. The product is recovered as the hydrochloride by dissolution in ethanol (40 ml), acidification with ether HCl and precipitation with ethyl ether (500 ml).

30 The white solid is filtered and dried under vacuum at 40°C.

Yield: 7.6 g (46.5%) - m.p.: 153-155°C

10

$[\alpha]^{589} (c = 1, \text{methanol}) = -13.5$

Example 2

a) Preparation of N-(2-tetralyl)-D-serine methyl ester

Sodium cyanoborohydride (0.07 moles, 4.5 g) is added to a solution of β -tetralone (0.068 moles, 10.5 g) and D-serine methyl ester hydrochloride (0.07 moles, 11 g) in 10/1 ethanol/methanol (550 ml). The mixture is reacted at r.t. for 24 hours, evaporated to dryness under vacuum, taken up with water (800 ml) and extracted with ethyl acetate (2 x 500 ml). The combined organic phases are extracted with 1N HCl (2 x 300 ml). The aqueous phases are alkalinized with sodium bicarbonate and extracted with ethyl acetate (3 x 200 ml). The combined organic phases are evaporated to dryness under vacuum, to yield the product as a yellow oil.

Yield: 12 g (72%)

b) Preparation of (R)-3-hydroxy-2-(1,2,3,4-tetrahydronaphthalen-2-(R,S)-ylamino)propanamide (CHF 2818)

Ammonia is bubbled through a solution of N-(2-tetralyl)-D-serine methyl ester (0.048 moles, 12 g) in methanol (150 ml), at 0°C, to a ~15M concentration. The hermetically sealed system is reacted at r.t. for 120 hours. The solution is evaporated to dryness under vacuum and the residual oil solidifies upon grinding in petroleum ether.

Yield: 9 g (80%) - m.p.: 104-115°C

c) Separation of CHF 2818 diastereomers

Preparation of 3-hydroxy-2-(R)-(1,2,3,4-tetrahydronaphthalen-2-(S)ylamino)propanamide (CHF 2983)

2-(R)-(1,2,3,4-tetrahydro-2-(R,S)-naphthalenylamino)-3-hydroxypropanamide (0.038 moles, 8.8 g) is

11

crystallized in ethyl acetate (200 ml) and the resulting solid is recrystallized in ethyl acetate (200 ml) twice and finally in ethanol (50 ml). The crystalline white solid is dried under vacuum at 45°C.

5 Yield: 1.9 g (yield 43%) - m.p. = 142-145°C

$[\alpha]^{589}$ (c = 1, methanol) = +92

d) Preparation of 3-hydroxy-2-(R)-(1,2,3,4-tetrahydro-naphthalen-2-(R)ylamino)propanamide hydrochloride (CHF 2982.01)

10 The mother liquors from the first three crystallizations of CHF 2983 are combined and left at 0°C for 48 hours: the precipitate is filtered off and the filtrate is evaporated to dryness under vacuum. The resulting wax solidifies by trituration in ethyl ether
15 (50 ml) at r.t.

The solid is recovered as the hydrochloride by dissolution in methanol HCl (3M, 20 ml) and evaporation to dryness under vacuum, then crystallized from 1/1 ethanol/ethyl acetate (400 ml). The crystalline white
20 solid is dried under vacuum at 45°C.

Yield: 1.8 g (36 %) - m.p. = 226 - 232°C

$[\alpha]^{589}$ (c = 1, methanol) = -105.1

Example 3

a) Preparation of 3-(4-methylphenyl)propanoyl chloride
25 3-(4-Methylphenyl)propanoic acid (0.055 moles, 9 g) is dissolved in thionyl chloride (1.008 moles, 120 g). The mixture is stirred for 30' at r.t., then refluxed for 1h 30', evaporated under vacuum to an oil and taken up with toluene and hexane, each time evaporating to
30 dryness.

Yield: 12.4 g

12

b) Preparation of 3-(4-methylphenyl)propanal

A solution of triphenylphosphine (0.117 moles, 30.8 g) in acetone (200 ml) is added under a nitrogen stream at r.t. with Cu (I) bis-(triphenyl-phosphine)-tetrahydroborate (0.067 moles, 40.69 g), then 3-(4-methylphenyl)propanoyl chloride (0.055 moles, 10 g) dissolved in acetone (85 ml) is dropped in 45'. The mixture is stirred at r.t. under nitrogen for 1h. The precipitated solid is filtered, washing with acetone and the filtrate is evaporated under vacuum. The residue is dissolved in chloroform (340 ml), added with cuprous chloride (0.135 moles, 13.38 g) and stirred under nitrogen stream for 1h at r.t. The mixture is filtered through celite, the filtrate is evaporated to dryness and the resulting residue is taken up into ethyl ether and petroleum ether, filtered and evaporated under vacuum to obtain an oil.

Yield: 6.7 g (83%)

c) Preparation of 3-hydroxy-2-(3-(4-methylphenyl)-propylamino)propanoic acid methyl ester hydrochloride

2% Sodium in methanol (0.045 moles, 51.7 ml) is added to D,L-Serine methyl ester hydrochloride (0.045 moles, 7 g), dissolved in methanol (70 ml), to obtain the free base.

The formed sodium chloride is precipitated with ethyl ether (150 ml) and filtered off. The filtrate is evaporated to dryness under vacuum. The resulting residue is dissolved in methanol (450 ml), added with 3-(4-methylphenylpropanal) (0.045 moles, 6.7 g) and adjusted to pH 6 with acetic acid, sodium

13

cyanoborohydride (0.048 moles, 3 g) is added and the mixture is reacted at r.t. for 24 hours. The mixture is acidified with methanol HCl, evaporated to dryness under vacuum, taken up with methylene chloride (600 ml), then
5 alkalized with triethylamine and washed with water (3 x 500 ml). The organic phase is evaporated to dryness under vacuum and the product is recovered as the hydrochloride taking up with ethyl ether (400 ml) and acidifying with ether HCl. The precipitated white solid
10 is dried under vacuum at 30°C.

Yield: 8.8 g (68%)

d) Preparation of 3-hydroxy-2-(3-(4-methylphenyl)propylamino)propanamide hydrochloride (CHF 2934.01)

3-Hydroxy-2-(3-(4-methylphenyl)propylamino)propanoic acid methyl ester hydrochloride (0.03 moles, 8,6 g)
15 is dissolved in water (500 ml), alkalized with 10% aqueous potassium carbonate to pH = 8. The free base is extracted with methylene chloride and evaporated to dryness under vacuum. The resulting pale yellow oil
20 (16.3 g) is dissolved in methanol (150 ml). Ammonia is bubbled through the solution, cooled at -5°C, to a ~15M concentration. The hermetically sealed system is reacted for 5 days at r.t., then evaporated to dryness under vacuum. The product is recovered as the
25 hydrochloride by dissolution in ethanol (40 ml), acidification with ether HCl and precipitation with ethyl ether (500 ml). The white solid is filtered and dried under vacuum at 40°C.

Yield: 4.9 g (60 %) - m.p. = 173-176°C

Example 4

a) Preparation of 4-phenylbutanoyl chloride

4-Phenyl-butanolic acid (0.83 moles, 13.57 g) is added to thionyl chloride (0.114 moles, 8.27 ml) and warmed to dissolve the solid. The mixture is stirred for 30' at r.t., then refluxed for 10', finally recovered as in example 3a).

A 100% yield is obtained (0.083 moles, 15.09 g).

b) Preparation of 4-phenylbutanal

The procedure as in 3b) is followed, starting from 4-phenylbutanoyl chloride (0.083 moles, 15.09 g), to obtain 11 g of product.

c) Preparation of (R)-3-hydroxy-2-(4-phenylbutyl-amino)propanoic acid methyl ester hydrochloride

2% Sodium in methanol (0.052 moles, 59.8 ml) is added to D-Serine methyl ester hydrochloride (0.052 moles, 8.16 g), dissolved in methanol (81.6 ml), to release the base. The formed sodium chloride is precipitated with ethyl ether (163 ml) and filtered off. The filtrate is evaporated to dryness under vacuum. The resulting residue is dissolved in methanol (150 ml), added with 4-phenylbutanal (0.051 moles, 10.69 g) and adjusted to pH 6 with acetic acid, added with sodium cyanoborohydride (0.055 moles, 3.62 g) and reacted at r.t. for 24 hours. The mixture is acidified with methanol HCl, evaporated to dryness under vacuum, taken up in methylene chloride (600 ml) and in a 1M sodium bicarbonate solution. The phases are separated and the aqueous one is extracted again with methylene chloride (3 x 200 ml), then the combined organic phases are washed with water and evaporated to dryness under vacuum. The product is

15

recovered as the hydrochloride, taking up with ethyl ether (180 ml) and acidifying with ether HCl. The precipitated white solid is dried under vacuum at 30°C.

Yield: 5.83 g (40%)

5 d) Preparation of (R)-3-hydroxy-2-(4-phenyl)butylamino)propanamide (CHF 2918)

(R)-3-(Hydroxy-2-(4-phenylbutylamino)propanoic acid methyl ester hydrochloride (0.02 moles, 5.83 g) is dissolved in water (250 ml) and alkalized with 10% aqueous sodium carbonate. The free base is extracted with methylene chloride (3 x 200 ml) and evaporated to dryness under vacuum. The resulting pale yellow oil is dissolved in methanol (150 ml). Ammonia is bubbled through the solution, cooled at -5°C, to a ~15M concentration. The hermetically sealed system is reacted for 5 days at r.t., then evaporated to dryness under vacuum to obtain a thick oil. The product is obtained as a solid taking up with ethyl ether (25 ml) and precipitating with hexane (400 ml). The white solid is filtered and dried under vacuum at 35°C.

15 Yield: 4.43 g (92 %) - m.p. = 59-61°C

With similar procedures as described in the examples 1 to 4 th compounds 1 to 5, 27, 28, 30, 33, 34, 38, 39, 42 to 44, 46 to 48, 52 and 53 of Table 1 were prepared.

25

Example 5

Preparation of (R)-3-hydroxy-2-(3-phenylpropylamino)propanamide (CHF 2679)

D-Serinamide (0.038 moles, 4 g) and 3-phenylpropanal (0.038 moles, 5.1 g) are dissolved in methanol (400 ml) in a Parr bottle and hydrogenated in

30

16

the presence of 10% Pd/C (3 g) at 40 psi, until hydrogen absorption ceases. The catalyst is filtered off and the solution is evaporated to dryness under vacuum, taken up in ethyl acetate (300 ml) and washed with water (2 x 200 ml). The organic phase is dried and dissolved in warm ethyl ether (300 ml) to precipitate the product as a white solid by slow evaporation of the solvent at r.t.

Yield: 3.8 g (45%) - m.p. = 76-78°C

$[\alpha]^{589}$ (c = 1, methanol) = +13.2°

With similar procedures as described in the example 5 the compounds 6 to 13, 15 to 22, 24, 26, 49 to 51 of Table 1 were prepared.

Example 6

Preparation of (R)-2-(4-heptylamino)-3-hydroxypropanamide phosphate (CHF 2870.02)

A solution of D-serinamide (0.01 moles, 1 g) and 4-heptanone (0.01 moles, 1.1 g) in methanol (50 ml) is added with 4M methanol HCl (0.0033 moles, 0.85 ml) and sodium cyanoborohydride (0.005 moles, 0.33 g). The mixture is reacted at r.t. for 10 days, acidified with methanol HCl to pH = 2, and evaporated to dryness under vacuum. The residue is taken up with water (100 ml), washed with ethyl ether (100 ml), alkalized with sodium carbonate and extracted with chloroform (3 x 100 ml). The resulting oil (1.3 g) is dissolved in methanol (50 ml) and treated with 85% phosphoric acid (0.0065 moles, 0.75 g), then evaporated to dryness under vacuum to obtain the product as a very light solid foam.

Yield: 2 g (77%) - m.p. = 150-156°C

$[\alpha]^{589}$ (c = 1, water) = +1.9

Example 7

a) Preparation of 3-hydroxy-2-(N-methyl-(3-phenylpropyl)amino)propanoic acid methyl ester

A solution of N-(3-phenylpropyl)serine methyl ester
5 (6.4 g, 0.027 mol) in methanol (150 ml) is added with
10% Pd-on-carbon (0.7 g) and 40% formic aldehyde (3.0 ml
0.04 mol) dissolved in methanol (50 ml). The mixture
is stirred at room temperature under hydrogen
atmosphere, under a slight pressure (40 psi) until
10 the absorption ceases. The mixture is filtered and
the filtrate is evaporated under vacuum. The residue
is taken up in ethyl ether (300 ml), washed with water
(2 x 200 ml), dried over sodium sulfate and
evaporated under vacuum. Yield: 6.6 g

b) Preparation of 3-hydroxy-2-(N-methyl-3-phenylpropyl)amino propanamide hydrochloride (CHF 2968.01)

The procedure of Example 3b is repeated.

Ammonia is bubbled through a solution of 3-hydroxy-
20 2-(N-methyl-(3-phenylpropylamino)propanoic acid methyl
ester (0.026 moles, 6.6 g) in methanol (150 ml), at 0°C,
to a ~ 15M concentration. The system is reacted at
T=80°C for 120 hours in a closed reactor. The solution
is evaporated to dryness under vacuum to obtain an oil
25 from which the product is separated by low pressure
liquid chromatography. The resulting oil is taken up in
absolute ethanol and ethyl acetate, then acidified with
ether HCl. The mixture is stirred, adding ethyl ether,
then the precipitate is filtered and dried under vacuum
30 at 40°C.

Yield: 3.5 g. - m.p. 112-114°C

With a similar procedure as described in the example 7 the following compounds were prepared:

(R)-3-hydroxy-2-(N-methyl-2-indanylamino)propanamide hydrochloride (CHF 3440.01; compound n. 63)

5 (S)-3-hydroxy-2-(N-methyl-2-indanylamino)propanamide hydrochloride (CHF 3462.01; compound n. 67).

Example 8

a) Preparation of methyl 2-(indanylamino)acetate

Glycine methyl ester (0.053 moles, 6.64 g) is
10 dissolved in absolute ethanol (420 ml) and methanol (42 ml), added with 2-indanone (0.053 moles, 7 g), and with sodium cyanoborohydride (0.058 moles, 3.7 g) under stirring, in 40 minutes. The mixture is left under stirring at r.t. overnight, evaporated to dryness under
15 vacuum and the residue is taken up with water (500 ml) and ethyl acetate (500 ml). The phases are separated and the organic phase is extracted with a 0.1N HCl solution (3 x 200 ml). The acidic solution is adjusted to pH = 8.5-9 with a sodium bicarbonate saturated solution and
20 extracted with ethyl acetate (3 x 250 ml). The organic solution is separated, dried over sodium sulfate and evaporated under vacuum. The resulting product is dried under vacuum at r.t.

Yield: 5.8 g (53.4%)

25 b) Preparation of 2-(2-indanylamino)acetamide hydrochloride (CHF 3381.01)

Methyl 2-(indanylamino)acetate (0.028 moles, 5.8 g) is dissolved in ~15M ammonia in methanol (150 ml) and kept in a closed test tube at r.t. for some days. The
30 solution is evaporated to dryness under vacuum, taken up with absolute ethanol (2 x 200 ml) and evaporated

19

each time. The oil is taken up into methanol (30 ml) and acidified to acid pH with a HCl solution in dry methanol under stirring. The product is precipitated by addition of ethyl ether, filtered and dried under vacuum at 45°C.

Yield: 6.05 g (94.6%) - m.p. = 212-213°C

Example 9

Preparation of 2-(N-methyl-2-indanylamino)acetamide (CHF 3488)

2-(2-indanylamino)acetamide (0.016 moles, 3.00 g) is dissolved in methanol (60 ml), and potassium carbonate (0.016 moles, 2.18 g) is added under stirring. Methyl iodide (0.028 moles, 4.14 g) is dropped therein at r.t. in 15 minutes. The mixture is reacted for 4 hours at r.t., then evaporated under vacuum. The resulting solid is taken up into water (100 ml), the aqueous solution is extracted with ethyl acetate (3 x 150 ml). The organic solution is dried over sodium sulfate, then evaporated under vacuum to obtain an oil which is chromatographed under medium pressure through silica gel (eluent chloroform:methanol = 90:10)

Yield: 1.31 g (40%) - m.p. = 122-123°C

Example 10

a) Preparation of methyl (S)-3-hydroxy-2-(2-indanylamino)propanoate

L-serine methyl ester hydrochloride (0.05 moles, 7.78 g) is dissolved in absolute ethanol (400 ml) and methanol (40 ml), added with 2-indanone (0.05 moles, 6.74 g) and sodium cyanoborohydride (0.55 moles, 3.64 g) under stirring in 30 minutes. The mixture is stirred at

20

r.t. for 5 hours, then evaporated to dryness under vacuum and the residue is taken up into water (150 ml) and ethyl acetate (150 ml). The phases are separated and the aqueous phase is further extracted with ethyl acetate (200 ml). The combined organic phases are extracted with a 0.2 N HCl solution (2 x 200 ml). The aqueous solution is separated, adjusted to pH = 8 with sodium bicarbonate and extracted with ethyl acetate (2 x 300 ml). The organic solution is separated, dried over sodium sulfate and evaporated under vacuum. The resulting product is dried under vacuum at r.t.

Yield: 8.4 g (71.4%)

b) Preparation of (S)-3-hydroxy-2-(2-indanylamino)-propanamide hydrochloride (CHF 2993.01)

Methyl (S)-3-hydroxy-2-(2-indanylamino)propanoate (0.0357 moles, 8.4 g) is dissolved in ~12M ammonia methanol (150 ml) and kept in a closed test tube at r.t. for some days. The solution is evaporated to dryness under vacuum to obtain an oil which is taken up into methanol (2 x 250 ml), evaporating to dryness each time. The resulting product (base) is taken up into ethyl acetate (170 ml) and acidified to acid pH with a HCl solution in 4.75N dry ethyl acetate, under stirring. The product is filtered, crystallized from absolute ethanol and dried under vacuum at 40°C.

Yield: 6.7 g (72.8%) - m.p. = 186-187°C

$[\alpha]^{589}$ (c=1, methanol) = +16.6 (hydrochloride)

$[\alpha]^{589}$ (c=1, methanol) = -24.6 (base)

With methods known in the art the mesylate salt (CHF 2993.02) was obtained.

21

m.p. 180-183°C

$[\alpha]^{589}$ (c = 1, methanol) = + 13.4

With a similar procedure as described in the example 10 the compounds 41, 57, 61, 62, 69, 72, 75 and 76 of Table 1 were prepared.

Example 11

Preparation of (S)-2-(2-indanylamino)-3-(2-methylpropanoyloxy) propanamide hydrochloride (CHF 3542.01)

(S)-3-hydroxy-2-(2-indanylamino)-propanamide hydrochloride (CHF 2993.01, 0.006 moles, 1.6 g) is dissolved in trifluoroacetic acid (3.75 ml) and added with 2-methylpropanoyl chloride (0.022 moles, 2.3 ml) by dropping. After 2 hours at r.t. the solution is evaporated under vacuum and the resulting oil is taken up into ethyl ether (100 ml). The product is filtered, ground in ethyl ether (50 ml) and recovered by filtration. The solid is dried under vacuum at r.t.

Yield: 1.8 g (90%) - m.p. = 171-174°C

With a similar procedure as described in the example 11 the following compounds were prepared:

(S)-3-acetyloxy-2-(3-phenylpropylamino)propanamide hydrochloride (CHF 3023.01; compound n. 45);

(S)-3-acetyloxy-2-(2-indanylamino)propanamide hydrochloride (CHF 3519.01; compound n. 74);

$[\alpha]^{589}$ (c = 1, methanol) = + 29.8

(S)-3-benzoyloxy-2-(2-indanylamino)propanamide hydrochloride (CHF 3548.01; compound n. 78)

30

Example 12

Preparation of (S)-3-hydroxy-2-(2-indanylamino)-N-methylpropanamide hydrochloride (CHF 3422.01)

N-(2-indanyl)-(S)-serine methyl ester (0.025 moles, 5.95 g) is dissolved in 8.03M methylamine solution in ethanol (155 ml) in a closed test tube. After 1 day the solution is evaporated under vacuum taking up the resulting oil with methanol (3 x 100 ml), petroleum ether (fract. 40-70, 100 ml) and evaporating each time. The resulting oil, taken up with petroleum ether (250 ml), solidifies under stirring. The filtered solid (~5.42 g) is dissolved in lukewarm ethyl acetate (250 ml) and added with HCl in dry ethyl acetate (3.5 M) under stirring until markedly acid pH. The product is filtered, washed repeatedly with ethyl ether (150 ml) and dried in oven under vacuum.

Yield: 5.33 g (77.9%) - m.p. = 190.5-192°C

$[\alpha]^{589}_{\text{DMSO}}$ (c=1) = +18

$[\alpha]^{589}_{\text{methanol}}$ (c=1) = -2

With a similar procedure as described in the example 12 the compounds 55, 56, 58, 60, 64 to 66 were prepared.

Example 13

a) Preparation of 5,6-dimethoxy-2-hydroxyimino-indanone

5,6-Dimethoxy-1-indanone (0.078 moles, 15 g) is dissolved in absolute ethanol (550 ml), thermostated at 50°C, and added with isoamyl nitrite (0.086 moles, 11.9 ml) and conc. HCl (11.9 ml). After a few minutes the product precipitates. The reaction is kept at 50°C for further 3 hours, then cooled at r.t. and the solid is

23

filtered washing with absolute ethanol (50 ml) and ethyl ether (100 ml). The product is dried in oven under vacuum at r.t.

Yield: 16.3 g (94.4%)

5 b) Preparation of 5,6-dimethoxy-2-indanylamine

 A solution of 5,6-dimethoxy-2-hydroxyimino-1-indanone (0.27 moles, 6 g) in glacial acetic acid (500 ml) is added with 96% sulfuric acid (3.3 ml) and 10% Pd/C (1.5 g). The mixture is hydrogenated in a Parr apparatus (r.t., 35 psi). When the hydrogen absorption ceases, the catalyst is filtered off through celite, washing with methanol (70 ml). The solution is evaporated to dryness, to obtain a white solid which is dissolved in water, then extracted with ethyl acetate (2 x 70 ml). The aqueous solution is alkalized with a 1M sodium hydroxide solution to pH = 8-8.5. The product is extracted with methylene chloride (2 x 70 ml). The organic solution is dried over sodium sulfate and evaporated under vacuum to obtain the solid product.

Yield: 4.6 g (88.5%)

c) Preparation of 2-(5,6-dimethoxy-2-indanylamino)-acetamide (CHF 3511)

 5,6-Dimethoxy-2-indanylamine (0.024 moles, 4.6 g) and sodium bicarbonate (0.026 moles, 2.2 g) are added to a solution of chloroacetamide (0.024 moles, 2.2 g) in absolute ethanol (100 ml) and refluxed for 10 hours. The mixture is filtered at r.t. and the solution is evaporated to dryness. The resulting oil is chromatographed under medium pressure through silica gel (eluent: methylene chloride/methanol = 90/10)

24

Yield: 1.95 g (32.7%) - m.p. = 135-138°C

With a similar procedure as described in the example 13 the following compound were prepared:

5 2-(5-fluoro-2-indanylamino)acetamide hydrochloride (CHF 3480.01);


2-(5,6-difluoro-2-indanylamino)acetamide hydrochloride (CHF 3518.01).

10 In the subsequent Table 1, the abbreviations and structure formulae of the compounds of the examples as well as those of other compounds obtained with the same methods as above described are reported.

Table 1

 α -Amino acids amides derivatives - Structure

5



R' is H, if not otherwise precised.

COMPOUND	R	R ₁	R ₂	STEREO- CHEM	COMP. N:
CHF 2043	(CH ₂) ₆ CH ₃	CH ₂ OH	H	RS	1
CHF 2088	(CH ₂) ₄ CH ₃	CH ₂ OH	H	R	2
CHF 2102	(CH ₂) ₆ CH ₃	CH ₂ OH	H	R	3
CHF 2452.01	(CH ₂) ₄ CH ₃	CH ₂ OH	H	S	4
CHF 2525.01	(CH ₂) ₆ CH ₃	CH ₂ OH	H	S	5
CHF 2545	CH(CH ₃) ₂	CH ₂ OH	H	R	6
CHF 2560	CH(CH ₃) ₂	CH ₂ OH	H	S	7
CHF 2571	(CH ₂) ₆ CH ₃	CH ₂ C ₆ H ₅	H	RS	8
CHF 2583	CH ₂ -cyclohexyl	CH ₂ OH	H	R	9
CHF 2597	(CH ₂) ₂ CH ₃	CH ₂ OH	H	S	10
CHF 2617	(CH ₂) ₆ CH ₃	CH ₂ OH	(CH ₂) ₆ CH ₃	R	11
CHF 2621.01	CH(CH ₃)(CH ₂) ₅ CH ₃	CH ₂ OH	H	R	12
CHF 2678	(CH ₂) ₂ C ₆ H ₅	CH ₂ OH	H	R	13
CHF 2679	(CH ₂) ₃ C ₆ H ₅	CH ₂ OH	H	R	14

- continued -

- continued -

COMPOUND	R	R ₁	R ₂	STEREOCHEM	COMP. N.
CHF 2721	(CH ₂) ₅ CH ₃	CH ₂ OH	H	RS	15
CHF 2722	(CH ₂) ₇ CH ₃	CH ₂ OH	H	RS	16
CHF 2723	(CH ₂) ₈ CH ₃	CH ₂ OH	H	RS	17
CHF 2739.01	$ \begin{array}{c} \text{CH}_2-\text{CH}-\text{CH}-\text{C}_2\text{H}_5 \\ \quad \\ \text{CH}_3 \text{CH}_3 \end{array} $	CH ₂ OH	H	RS	18
CHF 2750	$ \begin{array}{c} \text{CH}(\text{CH}_2)_5\text{CH}_3 \\ \\ \text{C}_2\text{H}_5 \end{array} $	CH ₂ OH	H	RS RS	19
CHF 2751	$ \begin{array}{c} \text{CH}(\text{CH}_2)_7\text{CH}_3 \\ \\ \text{CH}_3 \end{array} $	CH ₂ OH	H	RS RS	20
CHF 2768	1-(norbornyl- enyl)ethyl	CH ₂ OH	H	RS endo	21
CHF 2769	1-(norbornyl- enyl)ethyl	CH ₂ OH	H	RS exo	22
CHF 2803.01	(CH ₂) ₃ C ₆ H ₅	CH ₂ OH	H	S	23
CHF 2812	(CH ₂) ₃ C ₆ H ₅	CH ₂ OH	CH ₃	R	24
CHF 2818	1,2,3,4-tetra- hydro-2- naphthalenyl	CH ₂ OH	H	R RS	25
CHF 2824	$ \begin{array}{c} \text{CH}-(\text{CH}_2)_2\text{C}_6\text{H}_5 \\ \\ \text{CH}_3 \end{array} $	CH ₂ OH	H	R RS	26

- continued -

- continued -

COMPOUND	R	R ₁	R ₂	STEREOCHEM	COMP. N.
CHF 2847.01	$\begin{array}{c} \text{CH-CH}_2\text{C}_6\text{H}_5 \\ \\ \text{CH}_3 \end{array}$	CH ₂ OH	H	R S	27
CHF 2865.01	$\begin{array}{c} (\text{CH}_2)_2\text{-CHC}_6\text{H}_5 \\ \\ \text{CH}_3 \end{array}$	CH ₂ OH	H	R RS	28
CHF 2870.02	CH(CH ₂ CH ₂ CH ₃) ₂	CH ₂ OH	H	R	29
CHF 2880.01	$\begin{array}{c} \text{CH-C}_6\text{H}_5 \\ \\ \text{CH}_3 \end{array}$	CH ₂ OH	H	R RS	30
CHF 2918	(CH ₂) ₄ -C ₆ H ₅	CH ₂ OH	H	R	31
CHF 2934.01	3-(4-methyl-phenyl)propyl	CH ₂ OH	H	RS	32
CHF 2948.01	3-fluoro-phenyl)propyl	CH ₂ OH	H	RS	33
CHF 2967.01	3-(4-chloro-phenyl)propyl	CH ₂ OH	H	RS	34
CHF 2968.01	$\begin{array}{c} (\text{CH}_2)_3\text{-C}_6\text{H}_5 \\ \text{R}' = \text{CH}_3 \end{array}$	CH ₂ OH	H	RS	35
CHF 2982.01	1,2,3,4-tetrahydro-2-naphthalenyl	CH ₂ OH	H	R R	36
CHF 2983.01	"	CH ₂ OH	H	R S	37
CHF 2990.01	"	CH ₂ OH	H	S S	38
CHF 2991	"	CH ₂ OH	H	S R	39

- continued -

- continued -

COMPOUND	R	R ₁	R ₂	STEREOCHEM	COMP. N.
CHF 2993.01	2-indanyl	CH ₂ OH	H	S	40
CHF 2996	"	CH ₂ OH	H	R	41
CHF 3009.01	1,2,3,4-tetrahydro-2-naphthalenyl	CH ₃	H	S RS	42
CHF 3010.01	"	CH ₃	H	R RS	43
CHF 3011.01	2-naphthalenylmethyl	CH ₂ OH	H	RS	44
CHF 3023.01	(CH ₂) ₃ -C ₆ H ₅	CH ₂ OAc	H	S	45
CHF 3066	6-Meo-1,2,3,4-tetrahydro-2-naphthalenyl	CH ₂ OH	H	R RS	46

- continued -

- continued -

COMPOUND	R	R ₁	R ₂	STEREOCHEM	COMP. N.
CHF 3091.01	2-indanyl	CH ₃	H	RS	47
CHF 3107.01	3-(4-CF ₃ -phenyl)-propyl	CH ₂ OH	H	R	48
CHF 3145.01	1,2,3,4-tetrahydro-2-naphthalenyl	CH ₂ OH	C ₆ H ₅ CH ₂	R RS	49
CHF 3186.01	C ₆ H ₅ (CH ₂) ₃	CH ₂ OH	C ₆ H ₅ CH ₂	R	50
CHF 3189.01	C ₆ H ₅ (CH ₂) ₃	CH ₂ OH	CH ₃ (CH ₂) ₄	R	51
CHF 3195.01	C ₆ H ₅ (CH ₂) ₃ R' = C ₆ H ₅ -CH ₂	CH ₂ OH	H	R	52

- continued -

- continued -

COMPOUND	R	R ₁	R ₂	STEREOCHEM	COMP. N.
CHF 3293	(CH ₂) ₃ -C ₆ H ₅ R' = n-butyl	CH ₂ OH	H	R	53
CHF 3381.01	2-indanyl	H	H	-	54
CHF 3394.01	"	CH ₂ OH	CH ₃	R	55
CHF 3408.01	"	H	CH ₃	-	56
CHF 3414.01	1,2,3,4-tetra- hydro-2- naphthalenyl	H	H	RS	57
CHF 3421.01	2-indanyl	CH ₃	CH ₃	R	58
CHF 3422.01	"	CH ₂ OH	CH ₃	S	59
CHF 3427.01	"	CH ₃	CH ₃	S	60

- continued -

- continued -

COMPOUND	R	R ₁	R ₂	STEREOCHEM	COMP. N.
CHF 3431.01	2-indanyl	CH ₃	H	S	61
CHF 3434.01	"	CH ₃	H	R	62
CHF 3440.01	" R' = CH ₃	CH ₂ OH	H	R	63
CHF 3441.01	1,2,3,4-tetra- hydro-2- naphthalenyl	H	CH ₃	RS	64
CHF 3442.01	"	CH ₂ OH	CH ₃	R R	65
CHF 3443.01	"	CH ₂ OH	CH ₃	S R	66
CHF 3462.01	2-indanyl R' = CH ₃	CH ₂ OH	H	S	67
CHF 3480.01	5-F-2- indanyl	H	H	RS	68
CHF 3486.01	"	CH ₂ OH	H	S RS	69

- continued -

- continued -

COMPOUND	R	R ₁	R ₂	STEREOCHEM	COMP. N.
CHF 3488	2-indanyl R' = CH ₃	H	H	-	70
CHF 3511	5,6-MeO-2-indanyl	H	H	-	71
CHF 3512.01	"	CH ₂ OH	H	S	72
CHF 3518.01	5,6-F-2-indanyl	H	H	-	73
CHF 3519.01	2-indanyl	CH ₂ OAc	H	S	74
CHF 3531.01	5-MeO-2-indanyl	H	H	RS	75
CHF 3539.01	"	CH ₂ OH	H	S RS	76
CHF 3542.01	2-indanyl	CH ₂ OOC-iPr	H	S	77
CHF 3548.01	"	CH ₂ OOC-C ₆ H ₅	H	S	78

Anticonvulsant activity

The compounds of the invention were evaluated in some pharmacological tests in order to investigate their potential anticonvulsant activity. To this end, compounds were screened in the maximal electroshock (MES) test. This model is widely used to assess the efficacy of antiepileptic agents against generalized and partial seizures. This study was performed in both rat and mouse by using the experimental procedure described in W. Löscher et al., *Epilepsy Res.*, 2 (1988), 145-181. Briefly, a 60 Hz alternate current (mice 50 mA, rats 150 mA) was delivered for 0.2 sec through corneal electrodes by means of an electrical stimulator. Anticonvulsant potency of a compound was determined after 60 min and up to 180 min from administration (p.o.) by calculation of its ED₅₀ for suppression of tonic hind limb extensions. Groups of 10 animals per dose were used and the ED₅₀ was calculated from the dose-effect curve accordingly to Litchfield and Wilcoxon (*J. Pharmacol. Exp. Ther.*, 96 (1949), 99-113).

In the rat MES model only the time course of anticonvulsant activity was evaluated. Groups of 10 rats were treated at their equiactive dose 30, 60, 120, 240, 360 and 480 min before the establishment of the electroshock. The peak of anticonvulsant activity and the duration of action were then evaluated.

In another serie of experiments performed in mouse the neurotoxicity of the compounds was evaluated as a measure of the impaired motor function by using the horizontal screen test (L.L. Coughenour et al., *Pharmacol. Bioch. and Behav.*, 6 (1977), 351-353). In

this model mice are placed individually on top of a
squire wire screen which is mounted horizontally on a
metal rod, which is then rotated 180° so that mice are
on the bottom of the screens. Impairment of the motor
function is observed from the number of animals that
falls from the screen or fail to climb to the top of the
screens. The medial neurological toxic dose (TD₅₀) is
than calculated as above. The ratio of TD₅₀ and ED₅₀
refers to the therapeutic index (T.I.) for each
compound. The T.I. is used to show a useful separation
between neurotoxicity and antiepileptic activity. The
larger the T.I. would indicate a better separation
between the above activities and a good profile as
anticonvulsants.

In the mouse MES model all the examined compounds
showed a potency of anticonvulsant activity expressed in
moles better than that of milacemide and/or sodium
valproate. The ED₅₀ values were comprised between 1.2
and 0.5 mmol/kg with a potency ratio ranging from 4.5 to
35 with respect to milacemide and from 1 to 6.7 with
respect to sodium valproate.

The evaluation of the time course of anticonvulsant
activity showed that some compounds were rapidly
absorbed with a peak of the effect at 30 minutes after
administration, whereas other compounds exerted more
deferred effects that peaked even at three hours after
administration.

A common feature to many compounds was a good
persistance of the effects with a suppression of the
convulsions equal to or greater than 50% still
statistically significant more than three hours after

administration.

The duration of action was longer than that of milacemide which was of about one hour.

The results of the MES test in mouse of the more
5 representative compounds of the invention at 60 min
after administration are shown in the following
Table 2.

The activity of the compounds has been compared
with that of two compounds of the prior art: FCE 28245,
10 a prototype of a new serie of serine derivatives endowed
with an anticonvulsant activity and TEVA compound 2 (a
1-aminoindane derivative), described in WO 94/22808 and
in WO 95/18617 respectively.

15	Compound (CHF)	ED ₅₀ MES (mg/kg p.o.)	TD ₅₀ Horizontal screen (mg(kg p.o.)	T.I.
	2993	44	1172	27
	2996	34	> 1500	44
	2983	43	1300	30
20	2991	38	1099	29
	3431	22	251	11
	3440	35	689	20
	3381	21	274	13
	FCE 28245	180	1670	9
25	TEVA Comp. 2	38	299	8

T.I.: therapeutic index

All compounds were administered as HCl salts. The
individual enantiomers of 2-aminoindane and
30 aminotetraline derivatives showed a significant
anticonvulsant activity. The (S)-hydroxyl-2 aminoindane

derivative CHF2993 and its enantiomer (R) CHF2996 were equipotent in the MES test. This latter compound showed also the higher T.I. (44) since its low value of neurotoxicity (>1500 mg/kg p.o.). The introduction of a methyl-group in the moiety of the CHF2993 resulted in a compound (CHF3440) with the same anticonvulsant activity but with an increase of the neurotoxicity ($TD_{50}=689$). Conversely, both the (S) 2-(2-indanylamino) propionamide and the 2-(2-indanylamino) acetamide derivatives CHF3431 and CHF3381 were more potent in inhibiting MES but resulted more neurotoxic than the previously mentioned compounds. A good profile of anticonvulsant activity was also observed with the 3-hydroxy-2-tetralinamine derivatives CHF2983 (R,S enantiomer) and CHF2991 (S,R enantiomer).

All the tested compounds were about 3-4 times more potent than FCE28245, chemically 3-hydroxy-2-(4-(3-phenylpropyloxy)benzylamino)propanamide methanesulphonate.

It is also worth noting that the 2-aminoindane derivative CHF3381 had a higher T.I. than the prior art 1-aminoindane compound 2 known from WO 95/18617, chemically (S)-2-(indanylamino) acetamide. In fact the 2-aminoindane compound of the invention was more potent as an anticonvulsant than the 1-aminoindane of the prior art. Taken together these results it appears that the class of the 2-aminoindane compounds of the invention exhibited less impairment of the motor coordination than the compound of the prior art.

Further to the MES test in mouse, some compounds were also evaluated in the MES test in rat and in a

37

chemical model of tonic convulsions induced by bicuculline accordingly to the procedure described in Swinyard E.A. et al., Antiepileptic Drugs, 3rd Edition, Raven Press, New York (1989). In this model mice were
 5 observed for 30 minutes after administration of a dose of bicuculline (s.c.) that induced in 97% of the animals a presence of tonic convulsions. In animals treated with compounds, abolition of the hindleg tonic-extension component is taken as the end point thus suggesting that
 10 the substance under examination has the ability to prevent the seizure spread.

As reference compound FCE 26743 (S)-2-(4-(3-fluorobenzyloxy)benzylamino)propanamide was used, disclosed in the prior art in EP-B1-0400495.

15 The results are shown in Table 3 as follows:

Compound (CHF)	ED ₅₀ MES (MG/KG P.O.)	Peak of activity (hrs)*	Duration of action (hrs)**	ED ₅₀ bicucul- line mouse (mg(kg p.o))	
20 2993	31	3	6	65	
2996	32	1	2	n.t.	
2983	19	1	2	62	
FCE 26743	11	0.5	1	20	

25 *: time of the maximal effect

** : time at which the protection is statistically significant

n.t.: not tested

In the rat MES test CHF2993, CHF2996 and FCE 26743
 30 showed an anticonvulsant activity close to that found in

38

the same model in mouse. In this test CHF 2983 was more potent, with an ED₅₀ of 19 mg/kg p.o., lower than that reported in mouse MES. The analysis of the kinetics of the pharmacological effects showed that CHF2993 had the longest duration of action (6 hours). In any case the duration of action of all the tested compounds was longer than that of the reference compound FCE 26743 (1 hour). In the bicuculline-induced convulsions in mice, ED₅₀ for CHF2993, CHF2983 and FCE 26743 was 65, 62 and 20 mg/kg p.o., respectively. Although these values were higher than those obtained in mouse MES test, still these values were in the same order of potency of their MES anticonvulsant activity.

Taken together these results, we can conclude that the compounds described herein showed a significant anticonvulsant activity in the MES model in both mouse and rat and also in a model of bicuculline-induced convulsions in mouse. The activity exhibited by these compounds in mice were almost similar to that found for some standard antiepileptic drugs including phenytoin, carbamazepine, and at least four times higher than sodium valproate, compounds for which literature data evidentiate smaller therapeutic indexes in comparison with those found in the present investigation.

The compounds of the invention can be administered in a variety of dosage forms, e.g. orally, in the form of tablets, capsules, sugar or film coated tablets, liquid solutions; rectally, in the form of suppositories, parenterally, e.g. intramuscularly or by intravenous or injection or infusion. The therapeutic regimen for the different clinical syndromes must be

adapted to the type of pathology taking into account as usual, also the route of administration, the form in which the compound is administered and the age, weight and conditions of the subject involved.

5 In an embodiment the therapeutically effective amount is from about 1 mg to about 1000 mg, preferably from about 10 mg to about 300 mg.

Of course, these dosage regimens may be adjusted to provide the optimal therapeutic response.

10 The nature of pharmaceutical compositions containing the compounds of this invention in association with pharmaceutically acceptable carriers or diluents will, of course, depend upon the desired route of administration.

15 The compositions may be formulated in the conventional manner with the usual ingredients. For example, the compounds of the invention may be administered in the form of aqueous or oily solutions or suspensions, tablets, pills, gelatine capsules, syrups,
20 drops or suppositories.

Thus, for oral administration, the pharmaceutical compositions containing the compounds of this invention are preferably tablets, pills or gelatine which contain the active substance together with diluents, such as
25 lactose, dextrose, sucrose, mannitol, sorbitol, cellulose; lubricants, for instance silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; or they may also contain binders, such as starches, gelatine, methylcellulose,
30 carboxymethylcellulose, gum arabic, tragacanth, polyvinylpyrrolidone; disaggregating agents, such as

starches, alginic acid, alginates, sodium starch glycolate; effervescent mixtures; dyestuffs; sweeteners; wetting agents, such as lecithin, polysorbates, laurylsulphates; and, in general, non-toxic and pharmacologically inactive substances used in pharmaceutical formulations. Said pharmaceutical preparations may be manufactured in known manner, for example by means of mixing, granulating, tableting, sugar-coating, or film-coating processes.

The liquid dispersions for oral administration may be e.g. syrups, emulsions and suspensions.


The syrups may contain as carrier, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol. The suspension and the emulsions may contain as carrier, for example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol. The suspensions or solutions for intramuscular injections may contain together with the active compound a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g. propylene glycol, and if desired, a suitable amount of lidocaine hydrochloride.

The solutions for intravenous injections or infusion may contain as carrier, for example, sterile water or preferably they may be in the form of sterile aqueous isotonic saline solutions.

The suppositories may contain together with the active compound a pharmaceutically acceptable carrier, e.g. cocoa-butter, polyethylene glycol, a polyoxyethylene sorbitan fatty acid ester surfactant or lecithin.

CLAIMS

1. Compounds of formula I



wherein:

10 R is straight or branched alkyl; cycloalkylalkyl; arylalkyl or phenylalkyl optionally substituted at the ring with alkyl, halogen or haloalkyl; fused or non-fused aryl optionally substituted with alkyl, alkoxy, halogen or haloalkyl;

R' is hydrogen; alkyl; phenyl; phenylalkyl;

15 R₁ is optionally acylated C₁-C₄ hydroxyalkyl or phenylalkyl;

R₂ is hydrogen; alkyl; phenyl; phenylalkyl;

with the proviso that when R is aryl, R₁ can be hydrogen or alkyl C₁-C₄.

20 2. Compounds according to claim 1 wherein:

R₁ is CH₂OH; R is C₃-C₁₀ alkyl, phenyl-(C₂-C₄)-alkyl, or 1,2,3,4-tetrahydro-2-naphthalenyl or 2-indanyl optionally substituted with alkyl, alkoxy, halogen or haloalkyl; R' is hydrogen or methyl and R₂ is hydrogen.


25 3. Compounds according to claim 1 wherein R₁ is hydrogen or methyl, R is 1,2,3,4-tetrahydro-2-naphthalenyl or 2-indanyl optionally substituted with alkyl, alkoxy, halogen or haloalkyl and R' and R₂ are hydrogen.

30 4. Compounds according to claim 1 wherein R₁ is CH₂OH; R is phenyl-(C₂-C₃)-alkyl, 1,2,3,4-tetrahydro-2-

42

naphthalenyl, 2-indanyl; R' and R_2 are hydrogen.


5. A process for the preparation of the compounds of claims 1-4 which comprises reacting amino acids esters or amides of formula II



wherein R_1 is as defined above and X is an alkoxy group or a NHR_2 group, wherein R_2 is as defined above,
10 with compounds of formula III



wherein Y is an oxygen atom or a NH group, whereas R_3 and R_4 , which are the same or different, are hydrogen or, together with the carbon atom they are linked to, they form one of the groups R or R_1 , as defined above,
15 to give compounds of formula IV




which can then be transformed into compounds of formula I by means of one or more of the following reactions:

- 25 - when X is an alkoxy group, reaction with an amine of formula $\text{R}_2\text{-NH}_2$;
- N-alkylation;
- acylation of any hydroxy group present in R_1 ;
- salification and/or optical resolution.
30 - elimination of any protecting groups.

6. A process for the preparation of the compound of

43


claims 1-4 which comprises condensation of an α -halogen ester of formula V



5

wherein R_1 is as defined above and preferably H, W is a halogen atom (generally chlorine or bromine) and R_5 is an alkyl group, with an amine of formula VI

10



wherein R_3 and R_4 are as defined above, and subsequent amidation with the amine $\text{R}_2\text{-NH}_2$ wherein R_2 is as defined above to give compounds of formula (I).

15 7. Pharmaceutical compositions containing as the active ingredient a compound of claims 1-4 in admixture with suitable excipients or carriers.

8. The use of the compounds of claims 1-4 for the preparation of medicaments for the treatment of
20 neurodegenerative diseases.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/03773

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07C237/06 C07C237/20 A61K31/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PAOLO PEVARELLO ET AL. : "Reductive Alkylation of 'alpha'-Aminoamides" ORGANIC PREPARATIONS AND PROCEDURES INTERNATIONAL, vol. 28, no. 2, April 1996, pages 179-183, XP002045591 see page 179 - page 183; examples 3F,3G ---	1,2,4,7,8
X	US 5 332 840 A (SHARON D. REAL ET AL.) 26 July 1994 see example 8B ---	1,2
X	US 5 198 547 A (STEVEN W. BAILEY ET AL.) 30 March 1993 see column 14, line 45 - column 16, line 3 ---	1,2,5
X	EP 0 333 154 A (EISAI) 20 September 1989 see examples 2,43,44 ---	1,2
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

4 November 1997

Date of mailing of the international search report

11. 11. 97

Name and mailing address of the ISA
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Zervas, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/03773

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MARIA ALTAMURA ET AL.: "2-Substituted Penems with Amino Acid-Related Side Chains: Synthesis and Antibacterial Activity of a New Series of 'beta'-Lactam Antibiotics" JOURNAL OF MEDICINAL CHEMISTRY, vol. 38, no. 21, 13 October 1995, WASHINGTON US, pages 4244-4256, XP002045592 see page 4246, table 1, compounds 8f, 8g; page 4247, table 3, compounds 8f, 8g; page 4255, column 1, last paragraph - column 2, last paragraph ---	1,2
X	FRANCESCO DALLAVALLE ET AL.: "Complex Formation Equilibria of L-Amino-Acid Amides with Copper(II) in Aqueous Solution" HELVETICA CHIMICA ACTA, vol. 72, no. 7, 1 November 1989, BASEL CH, pages 1479-1486, XP002045593 see page 1485, line 12 - line 29 ---	1,5
X	GIANNI GALAVERNA ET AL.: "Chiral Molecular Laminates: Crystal Structures of Bis(N2-n-alkyl-(S)-phenylalaninamidato)copper(II) Complexes" TETRAHEDRON: ASYMMETRY., vol. 5, no. 7, July 1994, OXFORD GB, pages 1233-1240, XP002045594 see page 1234, line 18 - page 1235, line 7 ---	1,5
A	GB 2 048 852 A (CONTNENTAL PHARMA) 17 December 1980 cited in the application see claims; examples ---	1-8
A	WO 94 22809 A (PHARMACIA/FARMITALIA CARLO ERBA) 13 October 1994 cited in the application see the whole document ---	1-8
A	WO 94 22808 A (PHARMACIA/FARMITALIA CARLO ERBA) 13 October 1994 cited in the application see the whole document ---	1-8
A	EP 0 400 495 A (FARMITALIA CARLO ERBA) 5 December 1990 cited in the application see the whole document -----	1-8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 97/03773

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1,7,8
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
Claim 1 is so broad, that for determining the scope of a meaningful international search due, account has been taken of rule 33.3 PCT; special emphasis was put on on the examples during the international search.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/03773

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5332840 A	26-07-94	US 5436347 A	25-07-95
US 5198547 A	30-03-93	AU 3804193 A	21-10-93
		EP 0631581 A	04-01-95
		WO 9319069 A	30-09-93
		US 5350851 A	27-09-94
EP 333154 A	20-09-89	AT 155469 T	15-08-97
		CN 1036769 A,B	01-11-89
		DE 68928174 D	21-08-97
		EP 0612751 A	31-08-94
		EP 0764648 A	26-03-97
		GR 1000462 B	30-07-92
		JP 1308287 A	12-12-89
		JP 1991301 C	22-11-95
		JP 6099449 B	07-12-94
		JP 7179472 A	18-07-95
		PT 90022 B	31-05-94
		US 5128465 A	07-07-92
GB 2048852 A	17-12-80	LU 81068 A	08-10-80
		LU 81069 A	08-10-80
		AT 391134 B	27-08-90
		AT 392271 B	25-02-91
		AU 536499 B	10-05-84
		AU 5678480 A	25-09-80
		CA 1184567 A	26-03-85
		CH 645091 A	14-09-84
		DE 3010599 A	09-10-80
		DE 3050800 C	22-06-89
		DK 123580 A,B,	23-09-80
		FR 2451913 A	17-10-80
		JP 1462820 C	28-10-88
		JP 55143944 A	10-11-80
		JP 63009491 B	29-02-88
		NL 8001721 A,B,	24-09-80
		SE 453917 B	14-03-88
		SE 8002204 A	23-09-80
		US 4639468 A	27-01-87
		ZA 8001682 A	26-08-81

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/03773

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9422809 A	13-10-94	AT 152442 T	15-05-97
		AU 667169 B	07-03-96
		AU 6377694 A	24-10-94
		CA 2136008 A	13-10-94
		CN 1104018 A	21-06-95
		DE 69402935 D	05-06-97
		EP 0648202 A	19-04-95
		FI 945582 A	28-11-94
		HU 70944 A	28-11-95
		JP 8504828 T	28-05-96
		NZ 263183 A	27-02-96
		PL 306538 A	03-04-95
		US 5449692 A	12-09-95
		ZA 9401964 A	19-10-94

WO 9422808 A	13-10-94	AU 667164 B	07-03-96
		AU 6284294 A	24-10-94
		CA 2135783 A	13-10-94
		CN 1104017 A	21-06-95
		EP 0643688 A	22-03-95
		FI 945581 A	28-11-94
		HU 68256 A	28-06-95
		JP 7507814 T	31-08-95
		NZ 262907 A	27-02-96
		PL 306537 A	03-04-95
		US 5446066 A	29-08-95
		ZA 9401963 A	19-10-94

EP 400495 A	05-12-90	AU 645752 B	27-01-94
		AU 5729990 A	18-12-90
		CA 2033190 A	26-11-90
		CN 1047496 A,B	05-12-90
		CZ 9002520 A	12-06-96
		DE 69004337 D	09-12-93
		DE 69004337 T	24-02-94
		WO 9014334 A	29-11-90
		EP 0426816 A	15-05-91
		ES 2062174 T	16-12-94
		HU 9500703 A	28-12-95
		IE 63934 B	28-06-95

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/03773

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 400495 A		IL 94466 A	24-01-95
		JP 4500215 T	16-01-92
		NO 179944 B	07-10-96
		PT 94160 B	30-04-97
		US 5391577 A	21-02-95
		US 5502079 A	26-03-96
		US 5276611 A	04-01-94
		US 5236957 A	17-08-93
